



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

The ultimate solution, or at least the one that is most to be desired, it seems to me, will be a universal time scale which shall have its basis in paleontology and shall adhere to the classic names, and in which the cycles and epicycles of diastrophism will be regarded as probably the most useful criteria for delimiting formational or larger sedimentary units, but never *per se* as criteria for the division of geological time.

EDWARD W. BERRY

THE JOHNS HOPKINS UNIVERSITY

SPECIAL ARTICLES

THE INFLUENCE OF DRY VERSUS FRESH GREEN PLANT TISSUE ON CALCIUM ASSIMILATION

IN early work on mineral metabolism with both the cow and the goat we showed that milking animals, receiving grains and dry oat straw as a roughage, are brought into a decided negative calcium balance. In the case of a goat the interesting observation was made that after a period of negative calcium balance, followed by access to fresh green grass, a positive calcium balance was observed, using the same ration as was used in the period preceding the access to green plant tissue. In extensive experiments Forbes and associates and Meigs and his associates have observed negative calcium balances with milking cows receiving rations liberally supplied with calcium. The rations used were from air *dried* materials, supplemented in some cases with silage. The striking feature of all the data obtained in these experiments was the large amount of fecal calcium, indicating a failure to assimilate satisfactorily this base.

In these our preliminary experiments, we have used milking goats. They have readily been brought into negative calcium balance on a ration consisting of air dried grains and air dried straw, with more calcium excreted in the fecal residue than was ingested with the ration. When the *dry* cereal straw was displaced by an equivalent in dry matter of fresh *green* material, with no increase in the total calcium intake, the negative calcium

balance was reduced in one animal from 1.6–2.7 grams CaO to .6 CaO per day. With another animal it was reduced from 1.5–2.5 grams CaO per day to .3–.8 gram per day. On the low calcium intake of 8 to 9 grams of CaO per week we could not expect a positive calcium balance to ensue, but this remarkable difference in the amount of calcium assimilated from the two rations we believe, has very great significance.

These changes in calcium assimilation are not to be attributed to variation in water intake or to unavailability of the calcium. Apparently there is something having its source in fresh green materials, which controls or assists calcium assimilation. It is suggested that under the extra strain of rapid growth or milk production not enough calcium can be assimilated for the liberal uses made of this element, unless there is present an abundance of calcium in the diet as well as an abundance of this something that assists calcium assimilation. Possibly we are dealing with the anti-rachitic vitamine, assumed as the fourth food accessory factor. In any case this problem touches growth, milk production and egg production. In the case of nursing women the relation of diet to a positive or negative calcium balance and to dental conditions will assume new aspects.

The supposition that we are dealing with something influencing calcium assimilation and which is more abundant in green than in dried plant tissue and consequently variable with the season's milk, would explain the variations in the seasonal frequency of rickets, as observed and commented upon by Hess.

Our data are not yet inclusive enough to indicate definitely the factors involved in this problem, yet we have been sufficiently impressed with the constancy of the observations made that it appears desirable to re-emphasize this relation to mineral metabolism which we anticipated some years ago and expressed in an earlier publication.

E. B. HART,
H. STEENBOCK,
C. A. HOPPERT